THE UNIVERSITY OF TEXAS MDAnderson Cancer Center Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure

Making Cancer History®

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.

PRESENTATION **RECOMMENDED VACCINATIONS** • Administration of vaccines at least 14 days prior to elective procedure or chemotherapy is preferred • Otherwise administer at discharge or Day 14 (whichever is first) • May consider waiting at least 3 months (6 months if patient received Patient meets any of the following asplenic conditions: anti-CD20 monoclonal antibody, *i.e.*, rituximab) after completion of • Anatomic or splenectomy Provide the following chemotherapy/immunosuppressive regimens in order to maximize • Functional • Chronic graft-versus-host disease (cGVHD) vaccinations if not vaccine efficacy² • Post-splenic artery embolization previously administered¹ • Initial vaccinations³ \circ Splenic irradiation $\geq 50\%$ • Meningococcal - Meningococcal type B (Bexsero®) 0.5 mL IM and • Sickle cell disease - Meningococcal conjugate (Menveo[®]) 0.5 mL IM • Pneumococcal conjugate 20-valent (PCV20, Prevnar20[®])⁴ 0.5 mL IM • Haemophilus influenzae type b (Hib) 0.5 mL IM Follow-up vaccinations³ (long-term) Follow-up vaccinations³ (2 months after initial vaccination) • Meningococcal type B (Bexsero[®]) 0.5 mL IM • Booster 1 year after completion of the primary series and • Meningococcal • Meningococcal type B (Bexsero®) 0.5 mL IM and every 2-3 years thereafter • Meningococcal conjugate (Menveo[®]) 0.5 mL IM • Meningococcal conjugate vaccine (Menveo[®]) 0.5 mL IM • Booster 5 years from first dose and every 5 years thereafter

¹Only administer vaccines if not previously administered, except in hematopoietic stem cell transplant patients. Refer to Stem Cell Transplantation SOP GC2.6 Vaccination Guidelines for details. For non-stem cell transplant patients, if any part of a series has already been given, refer to follow-up vaccinations or Appendix A for further instructions. Refer to the Centers for Disease Control and Prevention (CDC) guidelines for comprehensive vaccination recommendations.

²Alternatively, may wait until immune reconstitution for patients receiving active chemotherapy. Revaccination outside of hematopoietic stem cell transplant patients is generally thought to be unnecessary but robust data is lacking. Antibiotic prophylaxis can be considered in addition to vaccination or for patients awaiting vaccination, see Appendix B.

³See Appendix C for guidance on administration of IM vaccines in patients at high-risk for bleeding

⁴Refer to Appendix A if patient had received pneumococcal conjugate 13-valent (PCV13) or pneumococcal polysaccharide (PPSV23). Refer to the CDC guidelines if patient had already received pneumococcal conjugate 15-valent (PCV15).

THE UNIVERSITY OF TEXAS MDAnderson Cancer Center Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patients

Making Cancer History®

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.

APPENDIX A: Vaccine Special Considerations¹

Meningococcal Vaccination	Pneumococcal Vaccination ²	<i>Haemophilus influenzae</i> type b (Hib) Vaccination
 Previously vaccinated with one dose of meningococcal conjugate (Menveo[®]) If ≤ 5 years since 1st dose, administer 2nd dose of series at least 8 weeks after and then administer booster every 5 years thereafter If ≥ 5 years past 1st dose, repeat 2 dose series and administer booster every 5 years thereafter Previously vaccinated with one dose of meningococcal type B (Bexsero[®]) If ≤ 3 years since 1st dose, administer 2nd dose of series at least 8 weeks after and then administer a booster 1 year later with subsequent boosters every 2-3 years thereafter If ≥ 3 years past 1st dose, repeat 2 dose series and administer a booster 1 year after completion of primary series with subsequent boosters every 2-3 years thereafter Meningococcal conjugate (Menveo[®]) can be given on the same day as meningococcal type B (Bexsero[®]) at different anatomical sites Meningococcal conjugate (Menveo[®]) and meningococcal type B (Bexsero[®]) should only be given IM (not subcutaneously) 	 Note: Booster doses of PCV20 (Prevnar 20[®]) are not necessary in patients who have already received a dose None or Unknown Administer 1 dose of PCV20 Previously vaccinated with only PPSV23 (Pneumovax 23[®]) Administer PCV20 at least 1 year after last PPSV23 dose Previously vaccinated with PCV13 (Prevnar 13[®]) but not PPSV23 Administer PCV20 at least 1 year after last PCV13 dose Previously vaccinated with PCV13 at any age and PPSV23 before age 65 years Administer PCV20 at least 5 years after last pneumococcal vaccine dose Previously vaccinated with PCV13 at any age and PPSV23 at age 65 years or older The decision to administer PCV20 at least ≥ 5 years after last pneumococcal vaccine dose is a shared clinical decision between the patient and the provider 	Booster doses of Hib are not necessary in patients who have already received a dose

PCV13 = pneumococcal 13-valent conjugate vaccine (Prevnar 13[®])

PCV20 = pneumococcal 20-valent conjugate vaccine (Prevnar $20^{\text{®}}$)

PPSV23 = pneumococcal polysaccharide 23-valent vaccine (Pneumovax 23[®])

¹Patients should receive other vaccines per CDC guidelines (*i.e.*, influenza yearly, Tdap every 10 years, *etc.*)

²Refer to the CDC pneumococcal vaccination summary or the CDC PneumoRecs VaxAdvisor clinical support tool for comprehensive pneumococcal vaccination recommendations

Page 2 of 7

THE UNIVERSITY OF TEXAS MDAnderson Cancer Center Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific particular to MD Ande

Page 3 of 7

Making Cancer History®

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.

APPENDIX B: Risk Factors for Infections with Encapsulated Organisms¹ and Relevant Prophylactic Agents

Patients should ideally be counseled on the below points prior to and after the procedure/chemotherapy. Family members of affected patients should ensure compliance with vaccines per CDC guidelines.

Travel considerations² and animal exposures:

- Visit health care provider or travel clinic 4-12 weeks prior to travel
- Care must be taken in areas where malaria is endemic (e.g., tropical regions such as sub-Saharan Africa) or where outbreaks of meningococcal disease occur
- Avoid tick bites and areas where babesiosis is endemic (e.g., upper Midwest United States and coastal New England)
- Animal bites should be immediately attended to, particularly dog bites (Capnocytophaga canimorsus)

Prophylaxis recommendations

Consider prophylaxis for high-risk patients:

- Hematologic malignancy patients with anatomic or functional asplenia/hyposplenia awaiting or with incomplete vaccination
- Prolonged prophylaxis independent of vaccination status may be pursued on a case-by-case basis:
- Hematopoietic stem cell transplant patients on therapeutic immunosuppressants for cGVHD
- For details refer to Stem Cell Transplantation SOP "GC2.6 Vaccination Guidelines"
- Patients with history of sepsis with encapsulated organism
- High-risk patients should be cautioned to seek medical attention for any febrile illness

Preferred regimens:

- Oral penicillin V potassium (250 mg 500 mg twice daily) or amoxicillin (500 mg twice daily)
- Second line cephalosporins (e.g., cephalexin 250 mg twice daily) are reasonable alternatives that may be tolerated by patients with mild non-IgE-mediated reactions to penicillin
- Third line fluoroquinolones (e.g., levofloxacin 500 mg daily) are less studied alternatives for patients with contraindications to penicillin or cephalosporins
- Can consider providing an emergency supply for high-risk patients not receiving prophylaxis to be used at onset of fever or infectious symptoms

¹ The most common vaccine-preventable infections in patients without a spleen are caused by the following encapsulated organisms: *Streptococcus pneumoniae*, *Neisseria meningitidis*, and *Haemophilus influenzae* ² Refer to CDC website on travelers' health for guidance on travel considerations and vaccine prophylaxis

THE UNIVERSITY OF TEXAS MDAnderson Cancer Center Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient pop

Making Cancer History®

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.

APPENDIX C: Administration of Vaccines to Patients with Increased Risk of Bleeding

Patients at risk of bleeding/hematoma from IM injections:

- Thrombocytopenic (platelets < 30 K/microliter)
- Receiving anticoagulation
- IM injections should be avoided in patients on warfarin with elevated INR (>4)
- Hemophilia
- Von Willebrand disease

Successful administration of certain IM vaccines has been accomplished in hemophilia patients with the following techniques:

- Use a 23 or 25-gauge needle (25-gauge is smaller and preferred if available) for vaccine administration
- Apply pressure to site of vaccination for two minutes afterwards
- \circ Do **not** rub or massage the site

CDC guidance:

- Patients on anticoagulation therapy presumably have a similar bleeding risk as patients with clotting disorders so providers should follow the same guidelines for IM administration
- When IM vaccine is indicated for a patient with bleeding disorder, the vaccine should be administered if physician familiar with patient's bleeding risk determines IM route is reasonably safe • Prior to administration of IM vaccine, patient and family should be informed of the risk of hematoma formation
- CDC discourages deviation from the recommended vaccine route, site, or dose for any vaccine since this can lead to reduced protection and increase the risk of an exaggerated local reaction

Page 4 of 7

THE UNIVERSITY OF TEXAS MDAnderson Cancer Center Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific pati

Page 5 of 7

Making Cancer History®

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.

SUGGESTED READINGS

- Bonanni, P., Grazzini, M., Niccolai, G., Paolini, D., Varone, O., Bartoloni, A., ... Bechini, A. (2017). Recommended vaccinations for asplenic and hyposplenic adult patients. *Human Vaccines & Immunotherapeutics*, 13(2), 359-368. https://doi.org/10.1080/21645515.2017.1264797
- Briere, E. C., Rubin, L., Moro, P. L., Cohn, A., Clark, T., & Messonnier, N. (2014). Prevention and control of haemophilus influenzae type b disease: Recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recommendations and Reports*, 63(1), 1-14. Retrieved from https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6301a1.htm

Carpenter, P. A., & Englund, J. A. (2016). How I vaccinate blood and marrow transplant recipients. *Blood*, 127(23), 2824-2832. https://doi.org/10.1182/blood-2015-12-550475

Centers for Disease Control and Prevention. (2024). Immunization Schedules. Retrieved from https://www.cdc.gov/vaccines/schedules/hcp/index.html

Centers for Disease Control and Prevention. (2024). Pneumococcal Vaccine Recommendations. Retrieved from https://www2a.cdc.gov/vaccines/m/pneumo/pneumo.html

Centers for Disease Control and Prevention. (2024). *Pneumococcal Vaccination: Summary of Who and When to Vaccinate*. Retrieved from https://www.cdc.gov/vaccines/vpd/pneumo/hcp/who-when-to-vaccinate.html

Centers for Disease Control and Prevention. (2024). Travelers' Health. Retrieved from https://wwwnc.cdc.gov/travel/

- Cohn, A. C., MacNeil, J. R., Clark, T. A., Ortega-Sanchez, I. R., Briere, E. Z., Meissner, H. C., ... Messonnier, N. E. (2013). Prevention and control of meningococcal disease: Recommendations of the Advisory Committee on Immunization Practices (ACIP). *Morbidity and Mortality Weekly Report. Recommendations and Reports*, *62*(2), 1-27. Retrieved from https://web.s.ebscohost.com/ehost/pdfviewer/pdfviewer?vid=0&sid=19d304fc-91a3-48af-9a65-2347a58a0c56%40redis
- Davies, J. M., Lewis, M. P. N., Wimperis, J., Rafi, I., Ladhani, S., & Bolton-Maggs, P. H. B. (2011). Review of guidelines for the prevention and treatment of infection in patients with an absent or dysfunctional spleen: Prepared on behalf of the British Committee for Standards in Haematology by a working party of the Haemato-Oncology Task Force. *British Journal of Haematology*, *155*(3), 308-317. https://doi.org/10.1111/j.1365-2141.2011.08843.x
- Freedman, M. S., Ault, K., & Bernstein, H. (2021). Advisory Committee on Immunization Practices recommended immunization schedule for adults aged 19 years or older-United States, 2021. *Morbidity and Mortality Weekly Report*, 70(6), 193-196. http://dx.doi.org/10.15585/mmwr.mm7006a2
- Halasa, N. B., Shankar, S. M., Talbot, T. R., Arbogast, P. G., Mitchel, E. F., Wang, W. C., ... Griffin, M. R. (2007). Incidence of invasive pneumococcal disease among individuals with sickle cell disease before and after the introduction of the pneumococcal conjugate vaccine. *Clinical Infectious Diseases*, 44(11), 1428-1433. https://doi.org/10.1086/516781
- Kanhutu, K., Jones, P., Cheng, A. C., Grannell, L., Best, E., & Spelman, D. (2017). Spleen Australia guidelines for the prevention of sepsis in patients with asplenia and hyposplenism in Australia and New Zealand. *Internal Medicine Journal*, 47(8), 848-855. https://doi.org/10.1111/imj.13348
- Kimberlin, D. W., Brady, M. T., Jackson, M. A. (Eds.). (2018). *Red Book (2018): Report of the Committee on Infectious Diseases*, 31st Edition. AAP publications. https://doi.org/10.1542/9781610021470
- Kobayashi, M., Pilishvili, T., Farrar, J. L., Leidner, A. J., Gierke, R., Prasad, N., ... Cohen, A. L. (2023). Pneumococcal vaccine for adults aged ≥19 years: Recommendations of the Advisory Committee on Immunization Practices, United States, 2023. *MMWR. Recommendations and Reports*, 72(3), 1-39. https://doi.org/10.15585/mmwr.rr7203a1

THE UNIVERSITY OF TEXAS MDAnderson Cancer Center Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient

Page 6 of 7

Making Cancer History®

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.

SUGGESTED READINGS - continued

- Rankine-Mullings, A. E., & Owusu-Ofori, S. (2017). Prophylactic antibiotics for preventing pneumococcal infection in children with sickle cell disease. *Cochrane Database of Systematic Reviews, 10*, CD003427. https://doi.org/10.1002/14651858.CD003427.pub4
- Rubin, L. G., Levin, M. J., Ljungman, P., Davies, E. G., Avery, R., Tomblyn, M., ... Kang, I. (2014). Executive summary: 2013 IDSA clinical practice guideline for vaccination of the immunocompromised host. *Clinical Infectious Diseases*, *58*(3), 309-318. https://doi.org/10.1093/cid/cit816
- Schiffer, C. A., Kari, B., Delaney, M., Hume, H., Magdalinksi, A. J., McCullough, J. J., ... Anderson, K. C. (2017). Platelet transfusion for patients with cancer: American Society of Clinical Oncology Clinical practice guideline update. *Journal of Clinical Oncology*, 36(3), 283-299. https://doi.org/10.1200/JCO.2017.76.1734

THE UNIVERSITY OF TEXAS MDAnderson Cancer Center Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patie

Making Cancer History®

Disclaimer: This algorithm has been developed for MD Anderson using a multidisciplinary approach considering circumstances particular to MD Anderson's specific patient population, services and structure, and clinical information. This is not intended to replace the independent medical or professional judgment of physicians or other health care providers in the context of individual clinical circumstances to determine a patient's care. This algorithm should not be used to treat pregnant women.

DEVELOPMENT CREDITS

This practice consensus statement is based on majority opinion of the Asplenia Management workgroup at the University of Texas MD Anderson Cancer Center for the patient population. These experts included:

Core Development Team Leads

Ella Ariza Heredia, MD (Infectious Diseases)

Workgroup Members

Karen Beaty, MSPA (Surgical Oncology) Jovan Borjan, PharmD (Pharmacy Clinical Programs) Natalie Dailey Garnes, MD (Infectious Diseases) Olga N. Fleckenstein, BS[•] Alison Gulbis, PharmD (Pharmacy Clinical Programs) Brittnee Macintyre, MSN, RN, FNP-C[•] Van Nguyen, PharmD (Pharmacy Clinical Programs) Kelsey Olson, PharmD (Pharmacy Clinical Programs) Amy Pai, PharmD[•] David Santos, MD (Surgical Oncology)

*Clinical Effectiveness Development Team

Page 7 of 7